



TAKHZYRO TREATMENT GOALS

Setting treatment goals with your doctor

Since hereditary angioedema (HAE) affects everyone differently and your treatment needs may change over time, it's important to regularly set treatment goals with your doctor.

Take a few moments to think about your current TAKHZYRO treatment plan. Then fill out this questionnaire and take it with you to your next doctor's appointment. You can use the space provided to write down answers and help you remember what was discussed.

Past treatment

Were you on a preventive treatment before taking TAKHZYRO?

- Yes No

If yes, for how long?

Current treatment

What about your TAKHZYRO treatment plan would you like to discuss with your doctor?

- Treatment effectiveness and safety
- Dosing and administration
- Other (please specify below)

WHAT IS TAKHZYRO?

TAKHZYRO is a prescription medicine used to prevent attacks of hereditary angioedema (HAE) in people 12 years of age and older.

It is not known if TAKHZYRO is safe and effective in children under 12 years of age.

IMPORTANT SAFETY INFORMATION

TAKHZYRO may cause serious side effects, including allergic reactions. Call your healthcare provider or get emergency help right away if you have any of the following symptoms:

- wheezing
- chest tightness
- faintness
- hives
- difficulty breathing
- fast heartbeat
- rash

Please see additional Important Safety Information throughout and full [Prescribing Information](#), including information for patients.

Recent treatment and attack history

How long have you been taking TAKHZYRO?

How many attacks have you experienced since starting TAKHZYRO? Please include all attacks you have experienced, even those you may consider minor.

0 1 2 3+

What was the average length of your attacks?

1-2 days 3-6 days 7+ days

Rate the average severity of your attacks:

Mild Moderate Severe

What areas of your body were affected by an attack? Check all that apply.

Face Throat Stomach
 Hands Genitals Feet
 Other

Have you been hospitalized in the past year due to an attack?

Yes, _____ time(s) No

Setting treatment goals (ask your doctor these questions at your next visit)

What are my treatment goals (eg, attack frequency, attack severity, other)?

What factors should I consider to stay on track with my treatment goals?

TAKHZYRO TIP:

Use the **TAKHZYRO Journey Journal** that came in your Welcome Kit to keep track of your doses, attacks, and progress on treatment goals.

IMPORTANT SAFETY INFORMATION (cont'd)

The most common side effects seen with TAKHZYRO were injection site reactions (pain, redness, and bruising), upper respiratory infection, and headache.

These are not all the possible side effects of TAKHZYRO. For more information, ask your healthcare provider or pharmacist. You may report side effects to the FDA at 1-800-FDA-1088.

TAKHZYRO has not been studied in pregnant or breastfeeding women. Talk to your healthcare provider about the risk of taking TAKHZYRO if you are pregnant, plan to be pregnant, are breastfeeding, or plan to breastfeed.

Please see additional Important Safety Information throughout and full [Prescribing Information](#), including information for patients.



©2021 Takeda Pharmaceutical Company Limited. 300 Shire Way, Lexington, MA 02421. 1-800-828-2088. All rights reserved. TAKEDA and the TAKEDA logo are trademarks or registered trademarks of Takeda Pharmaceutical Company Limited. TAKHZYRO is a registered trademark of Dyax Corp., a Takeda company. US-LANA-0700v1.0 01/21

Visit us on:



youtube.com/takhzyro
facebook.com/takhzyro

TAKHZYRO[®]
(lanadelumab-flyo) injection